

A deep *ROSAT* PSPC observation towards the CMB decrement close to PC1643+4631 A & B: no cluster X-ray emission

Rüdiger Kneissl, Rashid A. Sunyaev and Simon D.M. White

Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str.1, 85740 Garching, Germany

Accepted Received 17 December 1997

ABSTRACT

We report on a 16 ks observation with the *ROSAT* PSPC centred on the CMB decrement which Jones et al. (1997) report in the direction of the quasar pair PC1643+4631. We do not detect the X-ray emission which would be expected if the decrement were caused by Compton up-scattering of CMB photons by hot gas. Our upper limit on the bolometric X-ray flux within a circle of radius $1'5$ is 1.9×10^{-14} ergs cm $^{-2}$ s $^{-1}$ at 99.7% significance. This is based on data in the energy range 0.5–2.0 keV and assumes bremsstrahlung with an observed temperature of 2.5 keV. We investigate the requirements for a spherical body of hot gas to produce the decrement while remaining consistent with our data. With conservative assumptions we establish a lower redshift limit of 2.8 at 95 % confidence for a 10 keV isothermal object of any size. Even higher limits obtain for objects with temperatures between 0.2 and 10 keV. We conclude that the CMB decrement is unlikely to be caused by a galaxy cluster. We discuss alternatives to a single, spherical cluster, such as a double or elongated cluster, a supercluster/filament, or scattering by a moving cluster. None of these appears attractive, especially since another candidate for a decrement near a QSO pair and without an obvious associated cluster has been found. An explanation other than Comptonization by hot gas within a virialised object appears needed.

Key words: X-ray – cosmic microwave background – quasars:individual: PC1643+4631 A & B – galaxies:clusters

1 INTRODUCTION

In an intriguing paper Jones et al. (1997) report the discovery of a decrement in the cosmic microwave background (CMB) in the direction of the quasar pair PC1643+4631 A & B. They suggest this may be due to a distant galaxy cluster which might also produce the quasar pair through gravitational lensing of a single source. The lack of detected X-ray emission in this field enabled Jones et al to establish that any such cluster must be at a redshift of 1 or greater. In this paper we report results from a pointed *ROSAT* observation which improves this X-ray constraint by a factor of about 20.

The interferometric data of Jones et al were taken with the Ryle Telescope (RT) at 15 GHz and give a flux decrement of $-380 \pm 64 \mu\text{Jy}$ averaged over the $110'' \times 175''$ beam. For a circularly symmetric signal, this corresponds to a true decrement of at least 560 μK , this minimum value implying a core radius of about $60''$, or roughly $250h^{-1}$ kpc at high redshift. (Here $h = H_0/(100\text{km/s/Mpc})$.) Since intrinsic fluctuations in the CMB are not expected on arc-minute

scales, the most plausible origin for this decrement is inverse Compton scattering of CMB photons by hot intracluster gas (Sunyaev & Zel'dovich 1972). Decrements of this size have been detected in the direction of a number of massive galaxy clusters (Birkinshaw, Gull, Hardebeck 1984; Birkinshaw, Hughes, Arnaud 1991; Jones et al. 1993; Grainge et al. 1993; Birkinshaw, Hughes 1994; Wilbanks et al. 1994; Herbig et al. 1995; Grainge et al. 1996; Carlstrom, Joy, Grego 1997; Holzapfel et al. 1997; Myers et al. 1997). As Jones et al. discuss, such intracluster gas should be detectable in X-rays. They were able to put an upper limit on the X-ray flux using a 11.4 ks observation with the *ROSAT* PSPC in which the quasar pair serendipitously lies just within the field of view. Unfortunately vignetting and smearing by the point spread function greatly reduce the sensitivity of this observation, and we were able to establish a comparable (but independent) upper limit using data from *ROSAT* All-Sky Survey, which has a local exposure time of only 801 sec. Both observations limit the flux to be below 2×10^{-13} erg cm $^{-2}$ s $^{-1}$ (0.1–2.4 keV) at 99.7% confidence. Jones et al show that this constrains any cluster to be at $z > 1$. An extensive




Figure 1. Overlay of the CMB measurement (Jones et al. 1997) on our X-ray observation; coordinates in this plot are B1950. The separation of the quasar pair (+) PC1643+4631 A (right) & B (left) is $198''$. Positions of removed radio sources (x) are also marked. X-ray sources (see Tab. 1) are labelled 1–4.

search has so far identified no optical or infrared counterpart (Saunders et al. 1997). An HST observation has been scheduled (7342 by Saunders et al.) which should probe to even deeper limits.

If the observed decrement is indeed due to Compton scattering from hot gas, then X-ray observations offer the only way to detect this gas directly. The *ROSAT* detectors, with an energy range of 0.1–2.4 keV, are ideally suited to observe high redshift clusters which are expected to have temperatures in the range 1–10 keV. We therefore proposed and obtained a long pointed PSPC observation of this field.

2 THE *ROSAT* PSPC OBSERVATION

The observation, carried out between the 25th of February and the 3rd of March, was centred on RA = $16^{\text{h}}45^{\text{m}}12^{\text{s}}$, DEC = $+46^{\circ}24'35''$ (J 2000), and had a total accepted exposure time of 15889 s. This was among the very last observations with the PSPC before complete exhaustion of its gas supply. A small ($\sim 10\%$) drop in the gain occurred towards the end of our observation period and has been taken into account in the following analysis. Only few of the 61 X-ray sources detected in the observation have so far been identified, and all lie more than $35'$ off-axis. We were, nevertheless, able to check the absolute positioning of the *ROSAT* XRT by comparing several sources within the support structure with objects on the Palomar plates. This confirmed our estimated coordinates to be accurate to a few arcseconds. Our astrometry is thus much better than the width of the psf.

Fig. 1 shows an overlay of the CMB measurement on our X-ray map (in the 0.5 – 2.0 keV band). A pixelization 20 times finer than the local psf (which has FWHM of $37''.6$)

has been chosen, followed by Gaussian smoothing with a third of the width of the psf. The faintest grey patches are individual photons. We have labelled four sources near the decrement with a detection likelihood ratio exceeding 5, and we give their properties in Table 1. This very low detection threshold was chosen to demonstrate that not even a marginally detected source lies close to the centre of the microwave decrement. Secure detections are commonly quoted for a likelihood ratio of at least 10 ($\sim 4\sigma$). Objects 2 and 4 are both significantly more extended than the psf. Visual inspection suggests that 2 could consist of two close point sources, while 4 might really be an extended object. There are too few counts to use spectral analysis to check the possibility of a galaxy cluster. Other data (Brunner et al. 1996) suggest that X-ray emission from the quasars might be detectable in our data. We find no secure detection at the quasar positions, but there are marginal detections close to both of them (source 1 and possibly one component (with roughly half the total count rate) of the extended source 2). We use these to set 2σ upper limits of 7 and 20×10^{-4} ct s $^{-1}$ on sources associated with A and B respectively. For typical X-ray energy indices of $\alpha_x = 0.5$ –1.0 the monochromatic luminosities are 7.5 and 21×10^{44} erg s $^{-1}$ keV $^{-1}$, and with the optical magnitudes of $M_B = -26.1$ and -25.8 (Schneider, Schmidt, Gunn 1991) we find limits on the optical to X-ray spectral indices of $\alpha_{ox} > 1.4$ and 1.2.

The most stringent and robust upper limits on cluster X-ray emission associated with the decrement are obtained using the observed energy interval 0.5–2.0 keV, since the Galactic background is high at lower energies and the instrument has little sensitivity at higher energies. Results from including the 0.1–0.5 keV band, more stringent for intrinsic gas temperatures below 1 keV, are discussed later in the paper. We quote limits within a circular aperture of radius $1'.5$ centred on RA = $251^{\circ}29'58.3$, DEC = $+46^{\circ}40'68.7$, the centre of the decrement. This extends over the 3σ contours of the microwave observation. Within this area we have 13 counts compared to the expected background of 15 counts. (Consistent background estimates were obtained by a spline fit to the image after removal of detected sources, and by averaging random regions without obvious sources and more than $10'$ from the decrement.) With a Gaussian approximation to the Poisson photon arrival distribution we find that a source with an expected count of 12.5 can be excluded with 99.7% confidence. This corresponds to a limiting count rate of 8.9×10^{-4} cts s $^{-1}$. Deconvolving the effective area and response of the instrument, and correcting for Galactic absorption due to an HI column density of 1.8×10^{20} cm $^{-2}$, this corresponds to a bolometric flux limit of 1.9×10^{-14} erg cm $^{-2}$ s $^{-1}$ for an assumed bremsstrahlung spectrum with temperature 2.5 keV in the observer frame. This upper limit is almost twenty times lower than that found using the *ROSAT* Sky Survey data or the serendipitous observation analysed by Jones et al.

3 CLUSTER MODELS

We now study the constraints imposed on a possible high redshift cluster by the combined X-ray and radio data. We follow Jones et al. in modelling the intracluster medium as

Table 1. Source detections with likelihood > 5 near the centre of the decrement.

Object	RA (251°+)	DEC (46°+)	ML	Dist	Cts	Ext	Ext ML	Rate
1	.25658	.43541	6	132	4.9 ± 2.5			3.3 ± 1.7
2	.34073	.42180	73	122	43.8 ± 7.0	35	8	29.3 ± 4.7
3	.29930	.37334	18	152	10.6 ± 3.6			7.1 ± 2.4
4	.32700	.34584	78	263	64.1 ± 8.6	57	30	43.1 ± 5.8

ML: Likelihood ratio, Dist: distance to the centre of the CMB decrement in arcsec
 Cts: counts, Ext: extent (FWHM in arcsec), Rate: count rate in 10^{-4} ct s $^{-1}$




Figure 2. The amplitude of the true central temperature decrement (solid line) and the expected X-ray flux (in arbitrary units, dotted line) is plotted against the core radius for an isothermal model with density profile given by equ. 1. These data (points) were kindly provided by the Cambridge MRAO group.

an isothermal gas, temperature T , distributed spherically with density profile

$$n_e(r) = n_0 (1 + r^2/r_c^2)^{-1}. \quad (1)$$

To determine the central value of the temperature decrement a detailed knowledge of the synthesised beam, including the scanning path, is required. In practise the response of the telescope to different profiles must be modelled and fit directly to the visibility data. Jones et al. fit various core radii for the model of equ. 1, which implies a temperature decrement with profile

$$\Delta T = \Delta T_0 (1 + \Theta^2/\Theta_c^2)^{-\frac{1}{2}}. \quad (2)$$

The results were kindly provided to us by Richard Saunders and are shown in Fig. 2. For this model an X-ray flux limit F_x^{lim} within an aperture of radius Θ_a implies

$$(1+z)^4 r_A(H_0, \Omega, z) \frac{(kT)^2}{f(T)} > \frac{1}{4\pi} \left(\frac{m_e c^2}{\sigma T} \right)^2 \frac{\Theta_c y_0^2}{F_x^{lim}} \left(1 - [1 + \Theta_a^2/\Theta_c^2]^{-\frac{1}{2}} \right), \quad (3)$$

where $f(T) = \mathcal{L}_x/n_e^2$ is the X-ray cooling function, y_0 the central decrement, Θ_c the angular core radius of the cluster model, and $r_A(H_0, \Omega, z)$ is the angular-size distance. All quantities on the right of this inequality are directly observable, although in practice the radio data only constrain a combination of y_0 and Θ_c (Fig. 2). It turns out that a

Table 2. Gas halo parameters at $z \sim 1$ and 3 for $T = 10$ keV and the model of equ. 1

z	1.25	3	
Core radius	517	436	kpc
Central electron density	1.6	1.9	$\times 10^{-3}$ cm $^{-3}$
Emission integral	3.3	4.7	$\times 10^{-6}$ cm $^{-6}$ Mpc 3
X-ray luminosity (bol.)	2.0	2.8	$\times 10^{45}$ erg s $^{-1}$
X-ray limit (2- σ , bol.)	0.3	3.4	$\times 10^{45}$ erg s $^{-1}$

core radius near $1'$ not only yields the minimum true central temperature decrement, but also the lowest value of the right-hand side of equation 2 (for given F_x^{lim}). It thus gives the most conservative lower limit on the redshift of the cluster. Given the parameter dependence of the left-hand side, we will clearly get the lowest possible limit on $(1+z)$ if we take the largest plausible values for T and r_A . We will quote limits mostly for $T = 10$ keV, $h = 0.5$ and $\Omega = 1$. This choice for Ω is not conservative, but we will show below that our limits are only slightly reduced even for relatively extreme choices of the cosmological model.

The RT map is made for a very limited set of interferometer spacings, and hence only constrains the structure of the decrement on a small range of angular scales. As a result, interpretation of the radio data is quite model-dependent. The telescope response to model profiles other than that of equ. 2 has not been studied, so we present our limits for this case only. Since Θ_c is comparable to our X-ray resolution and also to the aperture we use to put limits on the X-ray flux, we do not expect a significant weakening of our limits for other reasonable profiles. (Equ. 1 is, in fact, a relatively good fit to most clusters studied so far.) We stress here that most plausible deviations from this model, for example a cooling flow in the inner regions, a temperature drop at large radii, or substructure in the gas, would actually enhance the X-ray emission relative to the microwave decrement, and thus would strengthen our limits.

Table 2 gives characteristic parameters for the gas halos at redshifts 1.25 and 3 which would produce the minimal X-ray emission consistent with the RT data; in both cases we assume $T = 10$ keV, $h = 0.5$ and $\Omega = 1$. For both redshifts the derived parameters are within the range observed for nearby clusters, although the luminosities are somewhat low and the core radii somewhat large as a result of our conservative assumptions. Table 2 also lists the upper limit on the X-ray luminosity implied by our PSPC observation. Clearly a cluster at $z = 1.25$ is strongly excluded, while a cluster at $z = 3$ is marginally consistent with our data. To explore the limits imposed on redshift and gas temperature in a more systematic way, we combine the radio and the X-ray data and maximise the joint likelihood of the two




Figure 3. The upper panel shows confidence contours for excluding the hot gas model of (equ. 1) based on the combined radio and X-ray data. The gas masses which yield the most consistent model at each redshift and temperature are shown in the lower panel (these are quoted in units of $10^{14} M_{\odot}$ and refer to the mass within a sphere of radius corresponding to $2r_c$). The heavy lines in the upper plot give confidence levels for an $\Omega = 1$ universe, while the dotted line gives the 95.4 % level for an open universe with $\Omega = 0.2$. The long and short dashed lines indicate the redshift of the quasar pair and of a damped Ly α absorption feature in the spectrum of quasar A. These plots are based on scalings derived from a thermal Bremsstrahlung model which assumes the standard helium abundance and no metals. We have checked the results against full spectral modelling including heavy elements, Galactic foreground absorption, and the response of the PSPC. Our results are little affected by these factors for emitted temperatures above 2 keV, but at lower temperatures they become sensitive to uncertainties in metallicity and absorption. Results from detailed spectral modelling for the allowed range of low gas temperatures are discussed in the text.

observations by varying the gas mass. Other model parameters are kept fixed, $h = 0.5$, $r_c = 50''$ and $\Omega = 1$ or 0.2. The results are presented in Fig. 3. Hot halos consistent with both datasets are required to lie at high redshift for any gas mass. For an assumed temperature of 5 keV, a redshift of 4.2 is required to get consistency at the 95% confidence level. This is beyond the redshift of the quasar pair ($z=3.8$). For an even hotter halo with $T = 10$ keV, the redshift is still required to exceed 2.8 at 95% confidence. As can be seen by comparing the limits for an Einstein-de Sitter universe and for an open universe with $\Omega = 0.2$, these limits are quite insensitive to the cosmological parameters. It is interesting that the hot cluster models which allow the lowest redshifts actually have rather small gas masses compared to nearby clusters of comparable temperature.

4 DISCUSSION

The X-ray flux predicted by our models could be reduced by a factor of 2 by assuming the cluster to be stretched along the line-of-sight to twice its apparent diameter or to be broken into two equal clusters which are superposed in projection. In fact, however, a factor of two is not a great help, since it only results in a ~ 25 % reduction of our redshift limits. If the decrement is due to a virialised cluster, it seems that we need a cluster at least as massive as Coma at a redshift beyond 3. The existence of even one such cluster in the observable universe is quite unexpected in current structure formation models. When normalised to give the presently observed abundance of clusters, only open models with $\Omega_0 < 0.25$ predict more than one cluster on the whole sky with $3 < z < 4$ and with $M > 2 \times 10^{15} M_{\odot}$. (This calculation is based on a Press-Schechter model similar to those in White et al (1993).) Such models overpredict the amplitude of fluctuations in the CMB.

Gravitationally confined gas which is considerably hotter than 10 keV has not been found observationally on the spatial scales of interest. At low temperature on the other hand large-scale gas properties are somewhat uncertain. The lower temperature window indicated by the drop in redshift below 2 keV in Fig. 3 might therefore be more promising for finding a plausible solution to the given data. Indeed the formal solution for a spherically symmetric object at redshift 3 and with temperature 0.2 keV could explain all observational data including the possible lensing for a gas mass fraction of 0.1. To derive this, the limit of $1.3 \times 10^{-3} \text{ ct s}^{-1}$ in the 0.1-2 keV energy interval was used, and very conservative assumptions about the plasma element abundances (zero metallicity, strongly suppressing line emission) and Galactic absorption ($N_H = 3.6 \times 10^{20} \text{ cm}^{-2}$, to account for local enhancements), which the limits in this temperature range depend sensitively upon, were made. The formation of such an object, however, would be hard to understand in the context of gravitational collapse. The implied gas mass of a few $\times 10^{15} M_{\odot}$ within a volume of $0.52h^{-3} \text{ Mpc}^3$ has not been seen in any object and the assumed temperature is by two orders of magnitude lower than what would be expected for the virial temperature.

Substantial reductions in X-ray flux are possible if we give up the assumption that we are seeing a collapsed and virialised system. The interferometer data require the extent of the object to be roughly $250h^{-1} \text{ kpc}$ in the directions transverse to the line-of-sight, but its depth could be much greater; it would then correspond to a supercluster “filament” seen end-on. Since such filaments are not fully virialised they can have temperatures which are substantially smaller than those of rich clusters, even though their masses are comparable or even larger. The gas in such filaments is heated by an accretion shock as it flows in from surrounding regions and by local virialisation shocks within subclumps along the filament. Typical velocities at these shocks are a few hundred km/s, so that post-shock temperatures of a few million degrees are expected. Thus at redshifts greater than 2 a prolate filament obeying equation (1) with $r_c = 50''$ but elongated by a factor of ten in depth (corresponding to a filament “length” of $\sim 5h^{-1} \text{ Mpc}$) can be consistent with both X-ray and radio data for gas temperatures below 0.5 keV. The gas mass required is around $1 \times 10^{15} M_{\odot}$.

(for $h = 0.5$). This high gas mass and the *a priori* implausible aspect of a close alignment of the filament with the line-of-sight disfavours this model, but careful quantitative analysis of numerical simulations is needed to see whether sufficiently massive filaments are formed at high redshift in currently popular theories for structure formation.

Gas in an aligned filament could also produce the observed microwave decrement without significant X-ray emission if it happened to be flowing along the filament at high enough speed. The amplitude of the observed decrement would be reproduced by Compton scattering from $3 \times 10^{14} M_{\odot}$ of gas moving away from us at 850 km/s, regardless of the gas temperature. The speed required scales inversely with the mass of gas, and, as before, the gas needs to be projected onto a region with transverse diameter about $0.5h^{-1}$ Mpc. Again numerical simulations are required to estimate whether sufficiently large coherent velocities occur along massive enough filaments for this to be a quantitatively viable model. Notice that this mechanism should produce enhancements and decrements of the CMB with equal probability.

The decrement towards PC1643+4631 may not be a unique, or even a particularly rare object. Another rather similar decrement has now been reported. A decrement with a smaller amplitude and on a smaller angular scale has been mapped at the VLA by (Richards et al. 1997). In this case also a deep *ROSAT* HRI observation found no X-ray emission. Furthermore, this decrement is also in the direction of an unusually close QSO pair at $z = 2.561$. We have applied our analysis to this case, for which $\Delta T_0 = -250 \mu\text{K}$, assuming $\Theta_c \sim 15''$ (comparable to the beam size), and the X-ray flux is less than $2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ at 99.7% confidence in the 0.1–2.4 keV band. For a 10 keV cluster these data imply a redshift greater than about 0.7. The constraint is weaker than in the case of PC1643+4631 because the decrement has smaller amplitude and scale. Thus PC1643+4631 may be the first example of a class of objects which is somehow related to the background QSO's. As Jones et al noted, the most obvious relationship might be through gravitational lensing, but to produce the observed $198''$ separation a filament at $z = 2$ would need to have a mass of $1 \times 10^{16} M_{\odot}$ projected between the two images. This seems very high, although again it cannot be excluded without detailed simulation of specific models.

5 CONCLUSION

Our X-ray limits make it seem unlikely that the decrement towards PC1643+4631 is due to a distant virialised galaxy cluster. We cannot definitively exclude the possibility of a hot, diffuse and massive cluster containing the two quasars at $z = 3.8$, but such a cluster is difficult to reconcile with present ideas about structure formation. A supercluster filament seen end-on at lower redshift may provide a more plausible solution, but then the association with the quasar pair is difficult to understand. It seems unlikely that this is pure chance since another similar decrement has been found also associated with a QSO pair. In view of these various problems it is undoubtedly important to constrain the microwave spectrum of the CMB decrement, to continue searching in other frequency ranges, as well as to explore other possi-

ble explanations, for example, topological defects such as textures or strings, exotic quasar phenomena, or foreground effects of various kinds.

ACKNOWLEDGEMENTS

We are very grateful to Joachim Trümper for supporting this *ROSAT* observation as part of the final PSPC operation programme, and we thank Hans Böhringer and Roland Egger for valuable suggestions throughout this work. We also thank Mike Jones and Richard Saunders for providing information on the RT's synthesised beam. RK wishes to thank Alastair Edge and Andy Fabian for helpful discussions, the Cambridge MRAO group and in particular Anthony Lasenby for their very kind hospitality during the completion of this paper and the SFB 375 Astro-Particle Physics for additional financial support.

REFERENCES

- Birkinshaw M., Gull S.F., Hardebeck H.E., 1984, *Nature* 309, 34
- Birkinshaw M., Hughes J.P., Arnaud K.A., 1991, *ApJ* 379, 466
- Birkinshaw M., Hughes J.P., 1994, *ApJ* 420, 33
- Brunner H., Lamer G., Friedrich P., Dörrer T., Stauber R., 1996, in: *Proc 'Röntgenstrahlung from the Universe'*, eds. H.U. Zimmermann, J. Trümper, H. Yorke, MPE Report 263, 433
- Carlstrom J.E., Joy M., Grego L., 1996, *ApJ* 456, L75
- Grainge K., Jones M., Pooley G., Saunders R., Edge A., 1993, *MNRAS* 265, L57
- Grainge K., Jones M., Pooley G., Saunders R., Baker J., Haynes T., Edge A., 1996, *MNRAS* 278, L17
- Herbig T., Lawrence C.R., Readhead A.C.S., Gulkis S., 1995, *ApJ* 449, L5
- Holzapfel W.L., et al., 1997, *ApJ* 480, 449
- Jones M., et al., 1993, *Nature* 365, 320
- Jones M.E., et al., 1997, *ApJ* 479, L5
- Myers S.T., Baker J.E., Readhead A.C.S., Herbig T., 1997, *ApJ* 485, 1
- Richards E.A., Fomalont E.B., Kellermann K.I., Partridge R.B., Windhorst R.A., 1997, *AJ* 113, 1475
- Saunders R., et al., 1997, *ApJ* 479, L1
- Schneider D.P., Schmidt M., Gunn J.E., 1991, *AJ*, 101, 2004
- Sunyaev R.A., Zel'dovich Ya., 1972, *Comments Astrophys. Space Phys.* 4, 173
- White S.D.M., Efstathiou G., Frenk C.S., 1993, *MNRAS* 262, 1023
- Wilbanks T.M., Ade P.A.R., Fischer M.L., Holzapfel W.L., Lange A.E., 1994, *ApJ* 427, L75

This paper has been produced using the Royal Astronomical Society/Blackwell Science $\LaTeX$ style file.